

March 2013

Dark matter scaling relations from mirror dark matter

R. Foot¹

*ARC Centre of Excellence for Particle Physics at the Terascale,
School of Physics, University of Melbourne,
Victoria 3010 Australia*

Mirror dark matter, and other similar dissipative dark matter candidates, need an energy source to stabilize dark matter halos in spiral galaxies. It has been suggested previously that ordinary supernovae can potentially supply the required energy. By matching the energy supplied to the halo from supernovae to that lost due to radiative cooling, we here derive an approximate scaling relation, $R_{SN} \propto \rho_0^2 r_0^3$ (R_{SN} is the supernova rate and ρ_0 , r_0 the dark matter central density and core radius). Such a relation is consistent with dark matter properties inferred from studies of spiral galaxies with halo masses larger than $3 \times 10^{11} M_\odot$. We also present a line of reasoning, related to the details of supernova energy transport in the halo, which explains why galaxies have a cored distribution and yields a second scaling relation, $r_0 \propto 1/\rho_0$. This relation is also in agreement with observations.

¹E-mail address: rfoot@unimelb.edu.au

A hidden sector exactly isomorphic to the ordinary matter sector is required if one hypothesizes fundamental improper space-time symmetries which are unbroken[1]. In such a theory, there is a mirror particle corresponding to every type of ordinary particle, except perhaps the graviton. Thus, a spectrum of stable dark particles naturally arises. We denote these mirror particles with a prime ('), $e', H', He', \dots$. The symmetry implies that the masses of these particles are identical to their ordinary matter counterparts and that the mirror particles interact with mirror gauge fields (such as the mirror photon) in a manner completely analogous to the ordinary matter sector.

Mirror particles have emerged as an interesting candidate for dark matter (for reviews and more complete bibliography see e.g.[2]). Mirror dark matter can explain[3] the positive dark matter signals from the DAMA[4], CoGeNT[5] and CRESST-II[6] direct detection experiments. This requires photon-mirror photon kinetic mixing (defined below) of strength $\epsilon \sim 10^{-9}$. Mirror dark matter can also explain[7] the large scale structure of the Universe (matter power spectrum and CMB) in a manner analogous to standard collisionless cold dark matter models provided[8] that $\epsilon \lesssim (1 - 3) \times 10^{-9}$.

On small scales mirror dark matter has a number of distinctive features due to self-interactions and dissipative interactions. In contrast to collisionless particles, galactic halos of spiral galaxies are composed predominately of mirror particles in a pressure supported spherical plasma[9]. There may also be a subcomponent consisting of compact objects such as old mirror stars[10] (gravitational lensing observations limit the MACHO halo fraction of the Milky Way to be less than around 0.3[11]). Because mirror dark matter is dissipative, an energy source is needed to stabilize the dark matter halos in galaxies. [Without an energy source the mirror particles would collapse to a dark disk on a time scale typically around a few hundred million years.] It has been speculated previously[9] that ordinary supernovae can potentially supply the required energy if photon - mirror photon kinetic mixing exists[12]:

$$\mathcal{L}_{mix} = \frac{\epsilon}{2} F^{\mu\nu} F'_{\mu\nu} \quad (1)$$

where $F_{\mu\nu}$ ($F'_{\mu\nu}$) is the field strength tensor for the photon (mirror photon). The physical effect of the kinetic mixing interaction[13] is to induce a tiny ordinary electric charge ($\propto \epsilon$) for the mirror charged particles. In the hot and dense core of type II supernovae mirror electrons and positrons can be produced from plasmon decay processes[14]. Thus ordinary supernovae can be a source of light mirror particles as well as the ordinary neutrinos. Indeed, it is estimated that half of the core collapse supernova energy is emitted by mirror particles ($e', \bar{e}', \gamma'$) if $\epsilon \sim 10^{-9}$ [14, 15]. A significant fraction of this energy might possibly be absorbed by the mirror particle halo. If this fraction is independent of galaxy size then an approximate scaling relation, $R_{SN} \propto \rho_0^2 r_0^3$ (R_{SN} is the supernova rate and ρ_0, r_0 the dark matter central density and core radius) follows by matching the energy supplied to the halo from supernovae to that lost due to radiative cooling. We find that this derived

relation is roughly consistent with dark matter properties inferred from studies of spiral galaxies. We also present a line of reasoning, related to the details of supernova energy transport in the halo, which potentially explains why galaxies, in this picture, have a cored distribution and yields a second scaling relation, $r_0 \propto 1/\rho_0$. This relation is also in agreement with observations. Although our discussion is in the context of mirror dark matter, similar conclusions could be obtained for a class of dissipative dark matter candidates which closely mimic mirror dark matter, such as those discussed recently in ref.[16].

The physical picture then, is that galactic halos in spiral galaxies are composed predominately of mirror particles $e', H', He', \dots$ in a self-interacting pressure supported halo. Clue's about the chemical composition of this halo arise from early Universe cosmology. Calculations indicate[17] a primordial mirror helium mass fraction around 0.9 for $\epsilon \sim 10^{-9}$. This suggests a halo composed primarily of He' with perhaps a fraction of H' and mirror metal components (produced in mirror star formation at an earlier epoch). To a first approximation, we can consider the halo as composed of mirror helium, which for temperatures above around 40 eV should be fully ionized. The radiative cooling rate, Γ_{cool} , of such a mirror particle plasma is given by the analogous expression for ordinary matter plasma[18]

$$\Gamma_{cool} = \Lambda(T) \int n_{e'}^2 dV \quad (2)$$

where $n_{e'}$ is the e' number density, and $\Lambda(T)$ is the cooling function and has units of $\text{erg cm}^3 \text{s}^{-1}$. For a mirror helium plasma, $n_{e'} \simeq 2\rho_{dm}/m_{He}$.

Rotation curves in spiral galaxies are well fit[19] with a baryonic component described by a Freeman disk[20] and a spherically distributed cored dark matter component with the Burkert profile[21]:

$$\rho_{dm} = \frac{\rho_0 r_0^3}{(r + r_0)(r^2 + r_0^2)} \quad (3)$$

where r_0 , ρ_0 is the dark matter core radius and central density respectively. However, as discussed in ref.[9], a spherical self gravitating isothermal gas of particles requires $\rho \propto 1/r^2$. If the temperature is not isothermal but rises in the inner region then it is possible to produce a cored distribution. This motivates the quasi-isothermal profile with $\rho_{dm} = \frac{\rho_0 r_0^2}{(r^2 + r_0^2)}$. Note that if, instead of the Burkert profile we adopted the quasi-isothermal profile, which also provides a reasonable fit to the rotation curves of spiral galaxies, this would not significantly affect our subsequent analysis. Anyway, assuming a mirror dark matter halo composed (predominately) of an ionized plasma with the Burkert density profile, it follows that

$$\Gamma_{cool} = \Lambda(T) \rho_0^2 r_0^3 (4/m_{He}^2) I \quad (4)$$

where

$$I \equiv 4\pi \int_0^\infty \frac{x^2}{(1+x)^2(1+x^2)^2} dx \simeq 1.34 . \quad (5)$$

Thus, we find:

$$\Gamma_{cool} \simeq \left(\frac{\Lambda(T)}{10^{-23} \text{ erg cm}^3/\text{s}} \right) \left(\frac{\rho_0 r_0}{10^{2.2} M_\odot/\text{pc}^2} \right)^2 \left(\frac{r_0}{10 \text{ kpc}} \right) 4 \times 10^{43} \text{ erg/s} . \quad (6)$$

On the other hand, the rate at which energy is supplied to the halo by ordinary core collapse supernovae is:

$$\begin{aligned} \Gamma_{SN} &= f_{SN} \langle E_{SN} \rangle R_{SN} \\ &\simeq \left(\frac{f_{SN}}{0.1} \right) \left(\frac{\langle E_{SN} \rangle}{3 \times 10^{53} \text{ erg}} \right) \left(\frac{R_{SN}}{0.03 \text{ yr}^{-1}} \right) 3 \times 10^{43} \text{ erg/s} \end{aligned} \quad (7)$$

where f_{SN} is the proportion of the supernova total energy, E_{SN} , supplied to the halo and R_{SN} is the galactic supernova rate. Equating Γ_{cool} with Γ_{SN} for the Milky Way galaxy implies $f_{SN} \sim 0.1$ with an order of magnitude uncertainty.

In this picture, $\Gamma_{cool} \simeq \Gamma_{SN}$ should hold for any galaxy, not just the Milky Way. Imposing this condition suggests a scaling relation:

$$R_{SN} \propto \Lambda(T) \rho_0^2 r_0^3 . \quad (8)$$

The idea is that the dynamics can keep this relation satisfied. If $\Gamma_{cool} > \Gamma_{SN}$ ($\Gamma_{cool} < \Gamma_{SN}$) then the halo should contract (expand), thereby increasing (decreasing) the star formation rate, and hence R_{SN} , until $\Gamma_{cool} \approx \Gamma_{SN}$. What is the temperature T ? For an isothermal halo in hydrostatic equilibrium, we expect[9]

$$T \approx \frac{1}{2} \bar{m} v_{rot}^2 \quad (9)$$

where $\bar{m}$ is the mean mass of the particles in the halo ($\bar{m} \approx 1 \text{ GeV}$ for a mirror helium dominated halo) and v_{rot} is the asymptotic value of the rotational velocity. For spiral galaxies, halo masses have values $3 \times 10^{10} M_\odot \lesssim M_h \lesssim 3 \times 10^{13} M_\odot$. For such halo masses, v_{rot} has the range $50 \text{ km/s} \lesssim v_{rot} \lesssim 500 \text{ km/s}$, and Eq.(9) suggests a rough temperature range:

$$10 \text{ eV} \lesssim T \lesssim 1000 \text{ eV} . \quad (10)$$

At the lowest temperature's mirror helium will not be fully ionized and the cooling rate can become dramatically suppressed. This might explain why low mass halos $< 10^{10} M_\odot$ hosting disk systems are not detected. For temperatures above $\sim 40 \text{ eV}$, mirror helium is fully ionized and for $40 \text{ eV} \lesssim T \lesssim 1000 \text{ eV}$, $\Lambda(T)$ typically varies by only a factor of 2-3 [18]. [This is relatively minor compared to the variation of, say, r_0 over this range of halo masses, which is around 2 orders of magnitude.] Taking the approximation of $\Lambda(T)$ as constant in this temperature range, yields the scaling relation:

$$R_{SN} \propto \rho_0^2 r_0^3 . \quad (11)$$

Is the above relation satisfied by spiral galaxies? A set of scaling relations have been derived from observations of rotation curves in spiral galaxies[22, 23, 24, 25] and summarized recently in ref.[26]. These relate ρ_0, r_0, M_h and r*-band luminosity, L_r :

$$\begin{aligned} \log \left(\frac{\rho_0 r_0}{M_\odot \text{pc}^{-2}} \right) &\simeq 2.2 \pm 0.25 \\ \frac{L_r}{1.2 \times 10^{10} L_\odot} &\simeq \frac{\left(\frac{M_h}{3 \times 10^{11} M_\odot} \right)^{2.65}}{1 + \left(\frac{M_h}{3 \times 10^{11} M_\odot} \right)^{2.00}} \\ \log \left(\frac{r_0}{\text{kpc}} \right) &\simeq 0.66 + 0.58 \log \left(\frac{M_h}{10^{11} M_\odot} \right) . \end{aligned} \quad (12)$$

These relations imply that for spiral galaxies with $M_h \gtrsim 3 \times 10^{11} M_\odot$,

$$L_r \propto r_0^{1.1} . \quad (13)$$

Observations[27] indicate that the galactic B-band luminosity, L_B , scales with the type II supernova rate as $L_B \propto R_{SN}^{1.3}$, with an uncertainty in the exponent around 0.1. Neglecting the expected minor difference between the scaling of L_B and L_r we arrive at an approximate ‘empirical’ scaling relation:

$$R_{SN} \propto r_0^{0.8} . \quad (14)$$

This relation is consistent, within the uncertainties, to the rough ‘theoretical’ scaling relation arrived at in Eq.(11) (given that $\rho_0 r_0$ is observed to be approximately constant). This provides some observational support to the notion that ordinary supernova supply the energy needed to stabilize halos in spiral galaxies, at least for $M_h \gtrsim 3 \times 10^{11} M_\odot$. What about for $M_h \lesssim 3 \times 10^{11} M_\odot$? These halos are expected to have a lower temperature $T \lesssim 40$ eV, for which the free mirror electron fraction reduces as e' become bound into atoms. This means that the right-hand side of Eq.(11) will begin to reduce more rapidly as a function of r_0 . This feature appears qualitatively consistent with observations as indicated by the sharper fall of the right-hand side of the second equation in Eq.(12) for $M_h \lesssim 3 \times 10^{11} M_\odot$.

The above analysis has assumed that a significant fraction ($f_{SN} \gtrsim 0.03$ or so) of supernova energy can be transmitted to the halo and that this fraction is roughly independent of galaxy size. How reasonable are these assumptions? Let us assume a kinetic mixing parameter $\epsilon \sim 10^{-9}$, so that around half of type II supernova energy is converted into $e', \bar{e}'$ emitted from the core with energies $\sim$ MeV. One could imagine that the huge number of MeV $e', \bar{e}'$ injected into a volume $[(\sim 1 \text{ pc})^3]$ around ordinary supernova will radiatively cool, converting most of their energy into mirror photons. The energy spectrum of these mirror photons is of course very hard to predict but it might be roughly analogous to the γ spectrum of ordinary Gamma Ray Bursts (GRB’s). GRB’s feature a wide spectrum of energies with mean around 700 keV with a few percent of energy radiated below 10 keV. In any case,

these mirror photons will then heat the mirror particle halo, potentially supplying the energy lost from the halo due to radiative cooling. Whether this can happen depends on how strongly the mirror photons scatter off mirror electrons, both bound and free.

Consider first the scattering off free mirror electrons, i.e. elastic (Thomson) scattering with E'_γ independent cross-section $\sigma_T = 6.7 \times 10^{-25} \text{ cm}^2$. We estimate that the optical depth due to elastic scattering for γ' propagating out from the galactic center is

$$\begin{aligned}\tau_{ES} &= \int_0^\infty \sigma_T n_{e'} dr \\ &\approx 0.78 \sigma_T \rho_0 r_0 \left(\frac{2}{m_{He}} \right) \sim 0.006\end{aligned}\quad (15)$$

where we obtained $\rho_0 r_0$ from Eq.(12). We expect $f_{SN} \sim \tau$ and Eq.(7) then suggests that elastic scattering is probably not frequent enough to supply enough heat to the halo to stabilize it. The cross-section is around an order of magnitude too small. However if the halo contains a significant proportion of heavy mirror elements - necessary to explain the direct detection experiments[3] - then the photoelectric cross-section of heavy mirror elements can easily dominate over the elastic cross-section for a large range of energies. This was noted in ref.[9] and we expand upon this point here.

Heavy elements, such as C', O', Si', Fe' , are not completely ionized but have their atomic inner shells filled. The total photoelectric cross-section (in units with $\hbar = c = 1$) for the inner K shell mirror electrons of a mirror element with atomic number, Z , is given approximately by[28]

$$\sigma_{PE}(E'_\gamma) = \frac{16\sqrt{2}\pi}{3m_e^2} \alpha^6 Z^5 \left[\frac{m_e}{E'_\gamma} \right]^{7/2}. \quad (16)$$

Evidently, the photoelectric cross-section decreases with mirror photon energy like $(E'_\gamma)^{-7/2}$ and, of course, E'_γ must be larger than the mirror electron binding energy of the particular element concerned. The contribution to the optical depth due to such inelastic scattering for γ' propagating out from the galactic center is

$$\begin{aligned}\tau_{IS} &= \sum_{A'} 2 \int_0^\infty \sigma_{PE} n_{A'} dr \\ &\sim \sum_{A'} 2 \rho_0 r_0 \sigma_{PE} \left[\frac{\xi_{A'}}{m_{A'}} \right]\end{aligned}\quad (17)$$

where $\xi_{A'}$ is the proportion by mass of the mirror metal component, A' (e.g. $A' = C', O', Si', Fe', \dots$) and we have included a factor of two since there are two K shell mirror electrons. For illustrative purposes we have evaluated the total optical depth, including both elastic and inelastic scattering (the latter assumed dominated by K shell bound electron scattering as discussed above) for an example with a 2% metal

component with $\xi_{C'} = \xi_{O'} = \xi_{Si'} = \xi_{Fe'} = 0.005$. The result is shown in figure 1. This figure indicates that for mirror photon energies

$$0.4 \text{ keV} \lesssim E'_\gamma \lesssim 30 \text{ keV} \quad (18)$$

inelastic scattering of mirror photons can dominate over elastic scattering. Thus, it might actually be possible for supernovae to transfer a significant part of their energy to the halo in a fairly efficient manner (i.e. $f_{SN} \gtrsim 0.03$).

Importantly, the optical depth is approximately independent of galaxy size since it scales as $\rho_0 r_0$ which is inferred to be constant [cf. Eq.(12)]. This means that the fraction of supernova energy transmitted to the halo is also roughly independent of galaxy size, which recall, was a necessary condition for the scaling relation, Eq.(11), to hold.




Figure 1: Galactic optical depth, τ , versus mirror photon energy (solid line). Also shown is the contribution to the optical depth due to elastic (Thomson) scattering, τ_{ES} (dotted line).

The scaling relation, Eq.(11), was derived from only the very general condition that the total energy lost due to radiative cooling is replaced by the total energy input. Of course in any particular volume element the energy input must match the energy output for a steady state configuration. It is useful to first consider a toy model, where we model the baryonic component as a point source whose energy output supports a spherical mirror dark matter halo. That is, we can assume a mirror photon luminosity, L , originating at $r = 0$. The energy going into a volume

element, $dV = 4\pi r^2 dr$, assuming mirror radiation dominates the energy transport is

$$\begin{aligned} d\mathcal{E}_{in} &= F\sigma ndV \\ &= \frac{L}{4\pi r^2}\sigma ndV \end{aligned} \quad (19)$$

where σ is the relevant mirror photon scattering cross-section [i.e. Thomson cross-section for $E'_\gamma \gtrsim 30$ keV and possibly a much larger photoelectric cross-section for E'_γ in the range given in Eq.(18)]. We have assumed in this toy model that the optical depth, $\tau \equiv \int \sigma n dr \ll 1$ so that the γ' flux, F , scales as $\sim 1/r^2$. The energy going out of the same volume element is

$$d\mathcal{E}_{out} = \Lambda(T)n^2 dV . \quad (20)$$

Matching $d\mathcal{E}_{in} = d\mathcal{E}_{out}$ implies

$$n = F \frac{\sigma}{\Lambda(T)} = \frac{L}{4\pi r^2} \frac{\sigma}{\Lambda(T)} . \quad (21)$$

The obtained $n \propto 1/r^2$ behaviour also matches the condition from hydrostatic equilibrium of a pressure supported self gravitating spherical distribution with a common (i.e. independent of r) temperature, T [9]. Thus a self consistent ‘toy model’ emerges, except it is unphysical at $r = 0$.

In the more realistic case there are two important perturbations to this picture. First the supernova energy sources are not all located at a point, $r = 0$, but a distribution extended over the disk scale length, r_D . This would be expected to soften the $F \propto 1/r^2$ behaviour at small $r \lesssim r_D$. Secondly, as figure 1 suggests, $\tau(E'_\gamma) \gtrsim 1$ for a range of E'_γ . This effect can increase the proportion of supernova energy being absorbed in the inner region of the halo versus the outer region. If this effect dominates, then the inner (core) region will heat up relative to the outer halo region. If the temperature of the core does indeed rise then the mirror metal components can have their inner shells completely ionized. If this happens then these elements can much less effectively absorb supernova energy. There will ultimately be a balance at some temperature $\sim T_c \lesssim E_b^{Fe}$. [Here $E_b^{Fe} \approx 8$ keV is the binding energy of the K shell mirror electrons for Fe' .] Thus we are lead to a picture where the inner region of the halo has temperature $T \sim T_c$ and the outer region has a somewhat lower temperature given by Eq.(9). The outer region should be roughly isothermal and we expect $\rho_{dm} \propto 1/r^2$, while the rise in temperature of the inner region will result in a much flatter or even decreasing dark matter density as $r \rightarrow 0$ (see discussion in the appendix).

The transition radius between the lower temperature outer halo and hotter inner halo, which roughly defines the core radius r_0 , will occur when $\langle \tau_{IS} \rangle \sim 1$. Here $\langle \tau_{IS} \rangle$ is the appropriate averaging of $\tau_{IS}(E'_\gamma)$ weighted by the γ' energy spectrum that supenovae ultimately produce. It might well be that the peak of this distribution is much greater than 10 keV (as indicated by the rough analogy with GRB’s mentioned

earlier), in which case the falling γ' spectrum in the relevant energy range $E'_\gamma \lesssim 30$ keV suggest that $\langle \tau_{IS} \rangle \sim \tau_{IS}(E'_\gamma \sim 10 \text{ keV})$. [Although the precise value of $\langle \tau_{IS} \rangle$ is uncertain, the important point is that it is not expected to scale significantly with galaxy size.] Thus, these arguments suggest that the existence of the core in galaxies is related to the heating up of the inner region, which in this picture, should occur when $\tau_{IS}(E'_\gamma \sim 10 \text{ keV}) \sim 1$ or

$$\begin{aligned} \sum_{A'} 2\rho_0 r_0 \left[\frac{\xi_{A'}}{m_{A'}} \right] \sigma_{PE}(E'_\gamma = 10 \text{ keV}) &\sim 1 \\ \Rightarrow \left(\frac{\rho_0 r_0}{10^{2.2} M_\odot \text{pc}^{-2}} \right) \left(\frac{\xi_{Fe'}}{0.02} \right) &\sim 1. \end{aligned} \quad (22)$$

The line of reasoning culminating in the above equation provides a simple physical explanation for the observed scaling: $r_0 \propto 1/\rho_0$, indicated in Eq.(12). Moreover, in this picture the observed value of $\rho_0 r_0$ provides, in principle, a measurement of $\xi_{Fe'}$ which it turns out is compatible with direct detection experiments[3] and CMB/LSS constraints[8].

In conclusion, we have considered galaxy structure within mirror dark matter - a dissipative and self-interacting dark matter candidate. For this type of dark matter, an energy source is needed to stabilize dark matter halos in spiral galaxies such as the Milky Way. Previously[9] it has been speculated that ordinary supernovae can supply the required energy if photon-mirror photon kinetic mixing of strength $\epsilon \sim 10^{-9}$ exists. We have shown here that this argument motivates an approximate scaling relation, $R_{SN} \propto \rho_0^2 r_0^3$ (R_{SN} is the supernova rate and ρ_0 , r_0 the dark matter central density and core radius). Interestingly, this scaling relation is consistent with the dark matter properties inferred from recent studies of spiral galaxies. We have also presented a line of reasoning, related to the details of supernova energy transport in the halo, which potentially explains why galaxies have a cored distribution and yields a second scaling relation, $r_0 \propto 1/\rho_0$. This relation is also in agreement with observations. Clearly more detailed studies are warranted to check the simple physical arguments presented here.

Appendix - hydrostatic equilibrium

Mirror dark matter, being self-interacting, forms a pressure supported halo. The pressure, P , can be determined from the hydrostatic equilibrium condition:

$$\begin{aligned} \frac{dP}{dr} &= -\rho(r)g(r) \\ &= -\bar{m}n(r)\frac{v_{rot}^2}{r} \end{aligned} \quad (23)$$

where $\rho(r) = \bar{m}n(r)$, with $\bar{m}$ being the mean mass of the mirror particles, e' , H' , He' , ..., in the plasma, $n(r)$ total dark matter particle number density, $g(r)$ is the local acceleration due to gravity. Here $P = nT$ and $v_{rot}(r)$ is the local rotational velocity.

For a spherically symmetric distribution, g (and hence v_{rot}^2/r) can be related to the total mass density, ρ_{total} , via:

$$g(r) = \frac{G}{r^2} \int_0^r \rho_{total} dV \quad (24)$$

As discussed in ref.[9], for an isothermal spherically symmetric halo, neglecting baryons, these equations have the solution $T = \frac{1}{2}\bar{m}v_{rot}^2$ and $\rho = \lambda/r^2$ ($\lambda \equiv \frac{T}{2\pi G\bar{m}}$). Of course, the approximation of neglecting baryons could only be roughly true for the outer regions of the halo.

Consider the case where the halo is not isothermal, but has a significantly hotter inner region. As briefly discussed already, this can be due to the efficient heating of the inner region due to the proximity of the supernova energy sources and significant opacity. The increasing temperature in the inner region means that the number density, $n(r)$, should be much flatter than $n(r) \propto 1/r^2$ (or even decreasing as $r \rightarrow 0$). If $n(r)$ were constant in the inner region, then integrating Eq.(23) implies

$$P(0) - P(r_0) \approx \bar{m}n \int_0^{r_0} \frac{v_{rot}^2}{r} dr \quad (25)$$

In the inner region of spiral galaxies, the rotational velocity, $v_{rot}(r)$, is roughly constant until very near the galactic center where $v_{rot} \rightarrow 0$ (see e.g.ref.[19]). It follows that

$$T(0) - T(r_0) \lesssim \bar{m}v_{rot}^2(r_0) \quad (26)$$

Given the isothermal result that T is related to the asymptotic v_{rot} value by: $T = \frac{1}{2}\bar{m}v_{rot}^2$, Eq.(26) indicates that a relatively modest increase in temperature in the inner region (less than a factor of two or so) would occur if $n(r)$ were constant there. As discussed in the text, it seems reasonable to expect the temperature of the central region to significantly increase until the mirror metal components are (almost) completely ionized, which suggests a central temperature around 1-10 keV. This is to be compared with the ‘average’ halo temperature, typically less than a keV, as indicated in Eq.(10). We thus anticipate that the temperature difference between the central region and outer halo region will, typically, be somewhat larger than that suggested in Eq.(26). If hydrostatic equilibrium were to hold then this suggests that the number density, $n(r)$, will likely be decreasing as $r \rightarrow 0$. Of course, this should not be a problem for observations (rotation curves), since the inner region of galaxies are well described by ordinary baryons[26].

Acknowledgments

The author would like to thank Shunsaku Horiuchi and Paolo Salucci for useful discussions/correspondence. This work was supported by the Australian Research Council.

References

- [1] R. Foot, H. Lew and R. R. Volkas, Phys. Lett. B **272**, 67 (1991); Mod. Phys. Lett. A **7**, 2567 (1992); R. Foot and R. R. Volkas, Phys. Rev. D **52**, 6595 (1995) [hep-ph/9505359]. Related concepts were discussed prior to the advent of the standard model in T. D. Lee and C. N. Yang, Phys. Rev. **104**, 256 (1956); I. Kobzarev, L. Okun, I. Pomeranchuk, Sov. J. Nucl. Phys. **3**, 837 (1966); M. Pavsic, Int. J. Theor. Phys. **9**, 229 (1974) [hep-ph/0105344].
- [2] A. Y. Ignatiev and R. R. Volkas, hep-ph/0306120; R. Foot, Int. J. Mod. Phys. D **13**, 2161 (2004) [astro-ph/0407623]; Int. J. Mod. Phys. A **19**, 3807 (2004) [astro-ph/0309330]; Z. Berezhiani, Int. J. Mod. Phys. A **19**, 3775 (2004) [hep-ph/0312335]; P. Ciarcelluti, Int. J. Mod. Phys. D **19**, 2151 (2010) [arXiv:1102.5530].
- [3] R. Foot, Phys. Rev. D **69**, 036001 (2004) [hep-ph/0308254]; Phys. Rev. D **78**, 043529 (2008) [arXiv:0804.4518]; Phys. Rev. D **86**, 023524 (2012) [arXiv:1203.2387].
- [4] R. Bernabei *et al.* (DAMA Collaboration), Riv. Nuovo Cimento. **26**, 1 (2003) [astro-ph/0307403]; Int. J. Mod. Phys. E **13**, 2127 (2004); Phys. Lett. B **480**, 23 (2000); Eur. Phys. J. C **67**, 39 (2010) [arXiv: 1002.1028]; Eur. Phys. J. C **56**, 333 (2008) [arXiv:0804.2741].
- [5] C. E. Aalseth *et al.* (CoGeNT Collaboration), Phys. Rev. Lett. **106**, 131301 (2011) [arXiv:1002.4703]; Phys. Rev. Lett. **107**, 141301 (2011) [arXiv: 1106.0650].
- [6] G. Angloher *et al.*, Eur. Phys. J. C **72**, 1971 (2012) [arXiv:1109.0702].
- [7] Z. Berezhiani, D. Comelli and F. L. Villante, Phys. Lett. B **503**, 362 (2001) [hep-ph/0008105]; A. Yu. Ignatiev and R. R. Volkas, Phys. Rev. D **68**, 023518 (2003) [hep-ph/0304260]; Z. Berezhiani, P. Ciarcelluti, D. Comelli and F. L. Villante, Int. J. Mod. Phys. D **14**, 107 (2005) [astro-ph/0312605]; P. Ciarcelluti, Int. J. Mod. Phys. D **14**, 187 (2005) [astro-ph/0409630]; Int. J. Mod. Phys. D **14**, 223 (2005) [astro-ph/0409633]. For pioneering work, see: S. I. Blinnikov and M. Yu. Khlopov, Sov. J. Nucl. Phys. **36**, 472 (1981); Sov. Astron. **27**, 371 (1983).
- [8] P. Ciarcelluti and R. Foot, Phys. Lett. B **679**, 278 (2009) [arXiv:0809.4438]; R. Foot, Phys. Lett. B **718**, 745 (2013) [arXiv:1208.6022].
- [9] R. Foot and R. R. Volkas, Phys. Rev. D **70**, 123508 (2004) [astro-ph/0407522].
- [10] Z. K. Silagadze, Phys. Atom. Nucl. **60**, 272 (1997) [Yad. Fiz. **60N2**, 336 (1997)] [hep-ph/9503481]; R. Foot, Phys. Lett. B **452**, 83 (1999) [astro-ph/9902065].

- [11] C. Alcock *et al.* [MACHO Collaboration], *Astrophys. J.* **542**, 281 (2000) [astro-ph/0001272]; P. Tisserand *et al.* [EROS-2 Collaboration], *Astron. Astrophys.* **469**, 387 (2007) [astro-ph/0607207].
- [12] R. Foot and X-G. He, *Phys. Lett. B* **267**, 509 (1991).
- [13] B. Holdom, *Phys. Lett. B* **166**, 196 (1986).
- [14] G. G. Raffelt, “Stars As Laboratories For Fundamental Physics: The Astrophysics Of Neutrinos, Axions, And Other Weakly Interacting Particles,” *Chicago, USA: Univ. Pr. (1996) 664 p.*
- [15] R. Foot and Z. K. Silagadze, *Int. J. Mod. Phys. D* **14**, 143 (2005) [astro-ph/0404515].
- [16] J. -W. Cui, H. -J. He, L. -C. Lu and F. -R. Yin, *Phys. Rev. D* **85**, 096003 (2012) [arXiv:1110.6893]; R. Foot, arXiv:1209.5602.
- [17] P. Ciarcelluti and R. Foot, *Phys. Lett. B* **690**, 462 (2010) [arXiv:1003.0880].
- [18] B. W. Carroll and D. A. Ostlie, “Modern Astrophysics,” *Addison-Wesley (1996).*
- [19] P. Salucci, A. Lapi, C. Tonini, G. Gentile, I. Yegorova and U. Klein, *Mon. Not. Roy. Astron. Soc.* **378**, 41 (2007) [astro-ph/0703115].
- [20] K. C. Freeman, *Astrophys. J.* **160**, 811 (1970).
- [21] A. Burkert, *IAU Symp.* **171**, 175 (1996) [*Astrophys. J.* **447**, L25 (1995)] [astro-ph/9504041]; P. Salucci and A. Burkert, *Astrophys. J.* **537**, L9 (2000) [astro-ph/0004397].
- [22] J. Kormendy and K. C. Freeman, astro-ph/0407321.
- [23] F. Donato and P. Salucci, *Mon. Not. Roy. Astron. Soc.* **353**, L17 (2004) [astro-ph/0403206].
- [24] M. Spano, *et al.*, *Mon. Not. Roy. Astron. Soc.* **383**, 297 (2008) [arXiv:0710.1345].
- [25] F. Shankar, A. Lapi, P. Salucci, G. De Zotti and L. Danese, *Astrophys. J.* **643**, 14 (2006) [astro-ph/0601577].
- [26] P. Salucci and M. De Laurentis, arXiv:1302.2268.
- [27] W. Li *et al.*, *Mon. Not. Roy. Astron. Soc.* **412**, 1473 (2011) [arXiv:1006.4613].
- [28] B. H. Bransden and C. J. Joachain, “Physics of Atoms and Molecules”, *Pren-tice Hall, 2nd Edition (2003).*